

AD-A042 734

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER  
ON ORTHOGONAL POLYNOMIALS. (U)

F/G 12/1

UNCLASSIFIED

JUN 77 P G NEVAI  
MRC-TSR-1760

DAAG29-75-C-0024  
NL

| OF |

ADA042 734



END  
DATE  
FILMED

9 - 77

DDC

12


ADA 042734

MRC TECHNICAL SUMMARY REPORT #1760  
ON ORTHOGONAL POLYNOMIALS  
Paul G. Nevai

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

June 1977

Received May 2, 1977



Approved for public release  
Distribution unlimited

DDC FILE COPY

Sponsored by

U. S. Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709

and

National Science Foundation  
Washington, D. C. 20550

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ON ORTHOGONAL POLYNOMIALS

Paul G. Nevai

Technical Summary Report #1760

June 1977

ABSTRACT

Orthogonal polynomials satisfy a three term recurrence relation. The purpose of the paper is to give estimates for the orthogonal polynomials and the corresponding weight function provided that the coefficients in the recurrence formula behave in a prescribed manner.

AMS(MOS) Subject Classification - 42A52

Key Words - Orthogonal Polynomials

Work Unit Number 6 - Spline Functions and Approximation Theory

EXPLANATION

Orthogonal polynomials provide a convenient means to expand functions into series in polynomials. When investigating these series one has to be able to estimate the size of the orthogonal polynomials. The present paper shows how to estimate orthogonal polynomials when the recurrence relation for these polynomials is given.

# ON ORTHOGONAL POLYNOMIALS

Paul G. Nevai

ACCESSION for	White Section	<input type="checkbox"/>
NTIS	Buff Section	<input type="checkbox"/>
DDC	Unannounced	<input type="checkbox"/>
USI Section		
DISTRIBUTION/AVAILABILITY CODES		
SP-CIAL		

Let  $\{\alpha\}_{n=0}^{\infty}$  and  $\{\gamma_n > 0\}_{n=0}^{\infty}$  be given sequences of real members. Putting  $p_{-1} = 0$ ,  $p_0 = \gamma_0$  and defining  $p_n$  for  $n = 1, 2, \dots$  by

$$xp_{n-1}(x) = \frac{\gamma_{n-1}}{\gamma_n} p_n(x) + \alpha_{n-1} p_{n-1}(x) + \frac{\gamma_{n-2}}{\gamma_{n-1}} p_{n-2}(x)$$

we obtain a system of polynomials  $\{p_n\}_{n=0}^{\infty}$  which by a result of J. Favard (see e.g. [2]) is orthonormal with respect to some positive measure  $d\alpha$  acting on the real line. Let

$$c_n = \left| 1 - 2 \frac{\gamma_{n-1}}{\gamma_n} \right| + 2|\alpha_{n-1}| + \left| 1 - 2 \frac{\gamma_{n-2}}{\gamma_{n-1}} \right|$$

It has been shown in [3] that under the assumption

$$(1) \quad \sum_{n=0}^{\infty} c_n < \infty$$

the measure  $d\alpha$  can be written as

$$d\alpha(x) = \alpha'(x)dx + \sum \{\text{jumps outside } (-1,1)\},$$

where  $\alpha'$  is positive and continuous on  $(-1,1)$  and  $\alpha'$  vanishes outside  $[-1,1]$ . At the present time it is not clear that assuming (1) how  $\alpha'$  behaves near  $-1$  and  $1$ . In case of the Tschebyshev polynomials ( $\alpha_n = 0$  for  $n = 0, 1, \dots$ ,  $\gamma_0 = \gamma_1 = 1$  and  $\gamma_n = 2^{n-1}$  for  $n = 2, 3, \dots$ )  $\alpha'$  is not continuous at  $-1$  and  $1$ . For the Tschebyshev polynomials of second kind ( $\alpha_n = 0$  and  $\gamma_n = 2^n$  for  $n = 0, 1, \dots$ )  $\alpha'$  is not positive at  $-1$  and  $1$ . Since the works of G. Szegő (see e.g. [4]) it has become known that those measures  $d\alpha$  for which

$$(2) \quad \int_{-\pi}^{\pi} \log \alpha'(\cos \theta) d\theta > -\infty$$

play a very important role in the theory of orthogonal polynomials. Therefore it is natural

Sponsored by

the United States Army under Contract No. DAAG29-75-C-0024 and  
the National Science Foundation under Grant No. MCS75-06687



to ask if (1) implies (2). It is easy to see that under the assumption  $\text{supp}(d\alpha) = [-1, 1]$  the inequality (2) follows from (1) ([3]). Otherwise the question is still open. It was proved in [3] that

$$(3) \quad \sum_{n=0}^{\infty} n c_n < \infty$$

implies

$$\alpha'(x) \geq \text{const} \sqrt{1-x^2}$$

for  $-1 \leq x \leq 1$ . Hence (2) follows from (3). K. M. Case suggested in [1] that (2) holds whenever

$$\limsup_{n \rightarrow \infty} n^2 c_n < \infty.$$

The purpose of this note is to show that the weaker condition

$$(4) \quad \sum_{n=0}^m (n+1) c_n \leq A \log(m+1) \quad (m = 1, 2, \dots)$$

not only implies (2) but also gives a pointwise estimate for  $\alpha'$ . We will see that assuming (4)  $\log \alpha'$  is very far from being nonintegrable. Our plan is the following. First, using an absolutely elementary method, we obtain estimate for  $|p_n|$ . This method is somewhat miraculous since we establish an inequality which improves itself when applied repeatedly. Having bound for  $|p_n|$  the corresponding estimate for  $\alpha'$  follows from a result in [3].

**THEOREM.** Suppose that (4) holds with a suitable constant  $A > 0$ . Then there exist two positive constants  $A_1$  and  $A_2$  depending only on  $A$  and  $\inf_n \gamma_{n-1}/\gamma_n$  such that

$$(5) \quad |p_n(x)| \leq A_1 (1-x^2)^{-A_2} \quad (-1 \leq x \leq 1)$$

for  $n = 1, 2, \dots$  and

$$(6) \quad \alpha'(x) \geq A_1^{-1} (1-x^2)^{A_2} \quad (-1 \leq x \leq 1).$$

**Proof.** Let  $x \in [-1, 1]$  and put  $x = \cos \theta$  where  $0 \leq \theta \leq \pi$ . Define  $\phi_n$  by

$$\phi_n(\theta) = p_n(x) - e^{i\theta} p_{n-1}(x).$$

Then

$$\phi_n(\theta) - e^{-i\theta} \phi_{n-1}(\theta) = p_n(x) - 2xp_{n-1}(x) + p_{n-2}(x)$$

and by the recurrence formula

$$\begin{aligned} (7) \quad \phi_n(\theta) - e^{-i\theta} \phi_{n-1}(\theta) &= \\ &= [1 - 2 \frac{\gamma_{n-1}}{\gamma_n}] p_n(x) - 2\alpha_{n-1} p_{n-1}(x) + [1 - 2 \frac{\gamma_{n-2}}{\gamma_{n-1}}] p_{n-2}(x). \end{aligned}$$

Consequently

$$|\phi_n(\theta) - e^{-i\theta} \phi_{n-1}(\theta)| \leq c_n \sum_{k=n-2}^n |p_k(x)|.$$

Using again the recurrence formula we obtain

$$(8) \quad \sum_{k=n-2}^n |p_k(x)| \leq K \sum_{k=M-1}^M |p_k(x)| \quad (M = n-1, n)$$

where  $K$  depends only on  $\sup_n \alpha_n$ ,  $\inf_n \gamma_{n-1}/\gamma_n$  and  $\sup_n \gamma_{n-1}/\gamma_n$ . Furthermore, from the definition of  $\phi_n$  follows that

$$(9) \quad \sqrt{1-x^2} |p_n(x)| \leq |\phi_n(\theta)|, \quad \sqrt{1-x^2} |p_{n-1}(x)| \leq |\phi_n(\theta)|.$$

Therefore

$$|\phi_n(\theta) - e^{-i\theta} \phi_{n-1}(\theta)| \leq 2Kc_n (1-x^2)^{\frac{1}{2}} \max_{|x| \leq 1} |\phi_{n-1}(\theta)|.$$

Recall that  $\phi_n - e^{-i\theta} \phi_{n-1}$  is a polynomial of degree  $n$  in  $x$ . Thus by a theorem of S. Bernstein

$$\max_{|x| \leq 1} |\phi_n(\theta) - e^{-i\theta} \phi_{n-1}(\theta)| \leq 2Kc_n (n+1) \max_{|x| \leq 1} |\phi_{n-1}(\theta)|,$$

that is

$$\max_{|x| \leq 1} |\phi_n(\theta)| \leq \max_{|x| \leq 1} |\phi_{n-1}(\theta)| [1 + 2Kc_n(n+1)] .$$

Repeated application of this inequality shows that

$$\max_{|x| \leq 1} |\phi_n(\theta)| \leq \gamma_0 \exp\{2K \sum_{j=1}^n (j+1)c_j\} .$$

Hence by (4)

$$(10) \quad |\phi_n(\theta)| \leq \gamma_0(n+1)^{2KA}$$

for  $-1 \leq x \leq 1$  and  $n = 0, 1, \dots$ . Now we return to (7). Multiplying both sides of (7) by  $e^{in\theta}$  and summing for  $n = 0, 1, \dots, m$  we obtain

$$\begin{aligned} e^{im\theta} \phi_m(\theta) &= \sum_{n=0}^m \left\{ \left[ 1 - 2 \frac{\gamma_{n-1}}{\gamma_n} \right] p_n(x) - 2\alpha_{n-1} p_{n-1}(x) + \right. \\ &\quad \left. + \left[ 1 - 2 \frac{\gamma_{n-2}}{\gamma_{n-1}} \right] p_{n-2}(x) \right\} . \end{aligned}$$

Therefore by (8) and (9)

$$(11) \quad |\phi_m(\theta)| \leq 2K(1-x^2)^{-\frac{1}{2}} \sum_{n=0}^m c_n |\phi_n(\theta)| .$$

Using inequality (10) we get

$$|\phi_m(\theta)| \leq 2K\gamma_0(1-x^2)^{-\frac{1}{2}} \sum_{n=0}^m c_n(n+1)^{2KA} .$$

If  $2KA < 1$  then by (4) and (9) the estimate (5) follows. Suppose that  $2KA > 1$ . Then using (4) we obtain

$$\begin{aligned} |\phi_m(\theta)| &\leq 2K\gamma_0(1-x^2)^{-\frac{1}{2}} (m+1)^{2KA-1} \sum_{n=0}^m c_n(n+1) \leq \\ &\leq 2KA\gamma_0(m+1)^{2KA-1} \log(m+1)(1-x^2)^{-\frac{1}{2}} \end{aligned}$$

which is much better than (10). Now plug this inequality into (11). If  $2KA - 1 < 1$  then (5) follows. Otherwise we get a new estimate which we again plug into (11). After finitely many similar steps we obtain



$$|\phi_m(\theta)| \leq B_1(1-x^2)^{-B_2}$$

for  $-1 \leq x \leq 1$  and  $n = 1, 2, \dots$  which combined with (9) yields (5). The inequality (6) follows from (5) and Theorem 7.5 of [3].

Finally we note that the example of Jacobi polynomials shows that apart from the constants  $A_1$  and  $A_2$  our result cannot be improved.

#### REFERENCES

- [1] Case, K. M., Orthogonal polynomials revisited, in "Theory and Application of Special Functions", ed. R. A. Askey, Academic Press, 1975, 289-304.
- [2] Freud, G., "Orthogonal Polynomials", Pergamon Press, New York, 1971.
- [3] Nevai, P. G., "Orthogonal Polynomials", Memoirs of the Amer. Math. Soc. (to appear).
- [4] Szegő, G., "Orthogonal Polynomials", Amer. Math. Soc., New York, 1967.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1760	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON ORTHOGONAL POLYNOMIALS.	5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Paul G. Nevai	8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024, MC975-06687	9. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit No. 6 - Spline Functions and Approximation Theory
10. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706	11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.	12. REPORT DATE June 1977
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 8p.	14. SECURITY CLASS. (of this report) UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 14 MRC-MSR-1760		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U.S. Army Research Office P.O. Box 12211 Research Triangle Park North Carolina 27709 and National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Orthogonal polynomials		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Orthogonal polynomials satisfy a three term recurrence relation. The purpose of the paper is to give estimates for the orthogonal polynomials and the corresponding weight function provided that the coefficients in the recurrence formula behave in a prescribed manner.		

221200

48